


## Peripheral Angioplasty / Stent for Atherosclerotic Occlusive Disease of the Lower Extremities 下肢動脈閉塞－血管成形(支架)手術

### Introduction

Atherosclerosis with build up of atherosclerotic plaques leads to narrowing or occlusion of the lower limb arteries. As a result, the supply of oxygen and nutrients to the lower limb becomes inadequate. It may cause leg pain during walking/ exercise that are usually relieved by rest, a condition known as intermittent claudication. In severe cases, rest pain, ulcer or gangrene may develop. In order to re-establish blood flow to the ischaemic part, surgical interventions may be required. Peripheral angioplasty/stenting may improve lower limb circulation.

### Procedure

The procedure is usually carried out under general, spinal or local anaesthesia. If the procedure is carried out under local anaesthesia, you will be awake throughout the procedure, but the anaesthetist will prescribe some sedatives for you. A small puncture is made at the groin. Under fluoroscopy guidance, a balloon catheter is used to dilate the narrowed or blocked vessel. A metallic stent may be placed into the artery if necessary. After successful angioplasty/stenting, patients usually stay in the hospital for 2 to 3 days.



## Risks & complications

- A. Perioperative mortality rate is usually less than 1 to 2%.
- B. Systemic complications
  - 1. Cardiovascular: myocardial infarction, arrhythmia, congestive heart failure
  - 2. Pulmonary: pneumonia, respiratory failure
  - 3. Renal failure/complication (contrast)
  - 4. Allergy/anaphylactic shock
  - 5. Others: cerebrovascular accident, deep vein thrombosis, pulmonary embolism
- C. Local complications
  - 1. Stent: infection, thrombosis
  - 2. Injury to the femoral arteries: haemorrhage, pseudoaneurysm
  - 3. Distal limb ischaemia
  - 4. Wound infection

## Before the procedure

- 1. You will be admitted one day before the procedure for preliminary tests including electrocardiogram, chest X-ray and blood tests.
- 2. The vascular surgeon will explain to you and your relatives about the procedure and its risks. You have to sign an informed consent.
- 3. Shaving of groin, shower / hair washing one day before the procedure.
- 4. Fasting of 6 hours prior to the procedure.
- 5. You have to wear a clean surgical gown, a cap and stockings immediately before the procedure.
- 6. Intravenous infusion, premedication of antibiotic may be required.

## Post-operative care

### General information

- 1. After general anaesthesia, you may feel tired, dizzy or weak. These will subside gradually.
- 2. Deep breathing and coughing exercise are helpful to prevent pneumonia.
- 3. We will monitor your blood pressure / pulse and observe your wound conditions.

### Activities

- 1. You should remain in bed for 12 hours. In particular, please do not move or bend the affected limb.
- 2. You may resume usual activities gradually under the instruction of the health care professional 12 hours after the procedure.
- 3. Avoid strenuous activities / heavy weight lifting within one week after the procedure.
- 4. Recovery depends on individual condition and progress.

### Wound care

1. Keep wound dressing clean and dry.
2. Observe the puncture site for haematoma, bleeding or oozing.
3. Take the prescribed analgesic as needed.

### Diet

1. Resumption of normal diet depends on the individual situation/ progress.
2. In general, you can take a well-balanced diet after the procedure unless you are on any special diet such as a diabetic or renal diet.

### Care after discharge

- A) Control the risk factors of atherosclerosis:
1. Quit smoking
  2. Go on low-fat, low-salt, low-sugar and high-fibre diet
  3. Take medications as prescribed to control hypertension / diabetes / high cholesterol level
- B) Proper foot care, avoiding any injury
- C) Maintain walking exercise regularly
- D) Drug therapy: continue drug therapy as prescribed, e.g. anti-platelet agent like aspirin
- E) Follow-up regularly with ultrasound imaging as arranged

### Remarks

This is general information only and the list of complications is not exhaustive. Other unforeseen complications may occasionally occur. In special patient groups, the actual risk may be different. For further information please contact your doctor.